

Phospho-IGF1R-Y1165/1166/INSR-Y1189/1190 Rabbit pAb

货号: B20836

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	154kDa/155kDa/156kDa
实测分子量	125kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa
细胞定位	cytoplasm,nucleus,plasma membrane,T-tubule
纯化	Affinity purification

抗原信息


抗原信息	A synthetic phosphorylated peptide around Y1165 & Y1166 of human IGF1RIGF1R (NP_000866.1).
序列	TDYYRK

靶点信息

研究背景	This receptor binds insulin-like growth factor with a high affinity. It has tyrosine kinase activity. The insulin-like growth factor I receptor plays a critical role in transformation events. Cleavage of the precursor generates alpha and beta subunits. It is highly overexpressed in most malignant tissues where it functions as an anti-apoptotic agent by enhancing cell survival. Alternatively spliced transcript variants encoding distinct isoforms have been found for this gene. [provided by RefSeq, May 2014]
基因ID	3480,3643

基因名	IGF1R,INSR
Swiss	P08069,P06213
别名	IGF1R/INSR

产品验证



Western blot analysis of Phospho-IGF1R-Y1165/1166/INSR-Y1189/1190 expressed in HeLa using Phospho-IGF1R-Y1165/1166/INSR-Y1189/1190 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn