

# PPP1R12B Rabbit pAb

货号: B20881

# 产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	
实测分子量	110kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,U-87MG
细胞定位	A band,cytoplasm,cytoskeleton,cytosol,nucleoplasm,Z disc
纯化	Affinity purification


# 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 660-930 of human PPP1 R12B (NP_002472.2).
序列	TFSRSRAERQAQEQPREKPTDTEGLEGSPEKHEPSAVPATEAGEGQQPWGRSLDEEPICHRLRCPAQPDKPTTPASPSTS RPSLYTSSHLLWTNRFSVPDSESSETTTNTTTAKEMDKNENEEADLDEQSSKRLSIRERRRPKERRRGTGINFWTKDEDET DGSEEVKETWHERLSRLESGGSNPTTSDSYGDRASARARREAREARLATLTSRVEEDSNRDYKKLYESALTENQKLKTKL QEAQLELADIKSKLEKVAQQKQEKTSDRSS

靶点信息

研究背景	Myosin phosphatase is a protein complex comprised of three subunits: a catalytic subunit (PP1c-delta, pr otein phosphatase 1, catalytic subunit delta), a large regulatory subunit (MYPT, myosin phosphatase targ et) and small regulatory subunit (sm-M20). Two isoforms of MYPT have been isolatedMYPT1 and MYPT2, the first of which is widely expressed, and the second of which may be specific to heart, skeletal muscle, and brain. Each of the MYPT isoforms functions to bind PP1c-delta and increase phosphatase activity. This locus encodes both MYTP2 and M20. Alternatively spliced transcript variants encoding different isoforms have been identified. Related pseudogenes have been defined on the Y chromosome. [provided by RefSe q, Oct 2011]
基因 <b>ID</b>	4660
基因名	PPP1R12B
Swiss	O60237
别名	MYPT2;PP1bp55;PPP1R12B

# 产品验证



Western blot analysis of PPP1R12B expressed in 293T,U-87MG using PPP1R12B Rabbit pAb at 1:1000. Se condary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

# 实验步骤

访问官网浏览详情: www.ablybio.cn