

SRPK2 Rabbit pAb

货号: B20900

产品信息

Human,Mouse,Rat
Rabbit
Polyclonal
WB IF/ICC
WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
120kDa
Liquid
Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
Unconjugated
Mouse heart,Mouse brain,Rat brain
cytoplasm,cytosol,nuclear speck,nucleolus,nucleoplasm,nucleus
Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 420-520 of human SRPK 2 (NP_872633.1).
序列	NAESDYTYSSSYEQFNGELPNGRHKIPESQFPEFSTSLFSGSLEPVACGSVLSEGSPLTEQEESSPSHDRSRTVSASSTGD LPKAKTRAADLLVNPLDPRN

靶点信息

研究背景	Serine/arginine-rich protein-specific kinase which specifically phosphorylates its substrates at serine resid ues located in regions rich in arginine/serine dipeptides, known as RS domains and is involved in the phos phorylation of SR splicing factors and the regulation of splicing. Promotes neuronal apoptosis by up-regul ating cyclin-D1 (CCND1 expression. This is done by the phosphorylation of SRSF2, leading to the suppress ion of p53/TP53 phosphorylation thereby relieving the repressive effect of p53/TP53 on cyclin-D1 (CCND1 expression. Phosphorylates ACIN1, and redistributes it from the nuclear speckles to the nucleoplasm, res ulting in cyclin A1 but not cyclin A2 up-regulation. Plays an essential role in spliceosomal B complex form ation via the phosphorylation of DDX23/PRP28. Probably by phosphorylating DDX23, leads to the suppres sion of incorrect R-loops formed during transcription; R-loops are composed of a DNA:RNA hybrid and the associated non-template single-stranded DNA. Can mediate hepatitis B virus (HBV core protein phosphorylation. Plays a negative role in the regulation of HBV replication through a mechanism not involving the phosphorylation of the core protein but by reducing the packaging efficiency of the pregenomic RNA (pgR NA without affecting the formation of the viral core particles.
基因 ID	6733
基因名	SRPK2
Swiss	P78362
别名	SFRSK2;SRPK2

产品验证



Western blot analysis of SRPK2 expressed in Mouse heart, Mouse brain, Rat brain using SRPK2 Rabbit pA b at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug pe r lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn