

TMF1 Rabbit pAb

货号: B20910

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	122kDa/123kDa
实测分子量	160kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	DU145,Raji,U-87MG
细胞定位	
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 900-1000 of human TMF1 (NP_009045.2).
序列	MKVEQERKKAIFTQETIKEKERKPFVSSTPTMSRSSSISGVDMAGLQTSFLSQDESHDSFGPMPISANGSNLYDAVRMGAGSSIIENLQSQLKLREGEI

靶点信息

研究背景	Potential coactivator of the androgen receptor. Mediates STAT3 degradation. May play critical roles in two RAB6-dependent retrograde transport processes: one from endosomes to the Golgi and the other from the Golgi to the ER. This protein binds the HIV-1 TATA element and inhibits transcriptional activation by the TATA-binding protein (TBP).
基因ID	7110

基因名	TMF1
Swiss	P82094
别名	ARA160;TMF;TMF1

产品验证



Western blot analysis of TMF1 expressed in DU145,Raji,U-87MG using TMF1 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn