

ELOVL2 Rabbit pAb

货号: B21044

产品信息

反应	Human,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	
实测分子量	38kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF7,HepG2,Rat testis
细胞定位	endoplasmic reticulum
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 247-296 of human ELOV L2 (NP_060240.3).
序列	ILFLNFYVQTYRKKPMKKDMQEPPAGKEVKNGFSKAYFTAANGVMNKKAQ


靶点信息

TIT 성숙되比	Ħ
御 先 自	H

Catalyzes the first and rate-limiting reaction of the four reactions that constitute the long-chain fatty acid s elongation cycle. This endoplasmic reticulum-bound enzymatic process allows the addition of 2 carbons to the chain of long- and very long-chain fatty acids (VLCFAs per cycle. Condensing enzyme that catalyze s the synthesis of polyunsaturated very long chain fatty acid (C20- and C22-PUFA, acting specifically toward polyunsaturated acyl-CoA with the higher activity toward C20:4(n-6 acyl-CoA. May participate in the production of polyunsaturated VLCFAs of different chain lengths that are involved in multiple biological processes as precursors of membrane lipids and lipid mediators.

基因ID	54898
基因名	ELOVL2
Swiss	Q9NXB9
别名	SSC2;ELOVL2

产品验证



Western blot analysis of ELOVL2 expressed in MCF7,HepG2,Rat testis using ELOVL2 Rabbit pAb at 1:10 00. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. B locking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn