

RANBP10 Rabbit pAb

货号: **B21056**

产品信息

反应	Human, Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	58kDa/64kDa/67kDa
实测分子量	73kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	U-87MG,HeLa,Mouse spleen
细胞定位	cytoplasm,cytosol,nucleus
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 362-497 of human RAN BP10 (NP_065901.1).
序列	YPGSPSLSPRHGPSSSHMHNTGADSPSCSNGVASTKSKQNHSKYPAPSSSSSSSSSSSSSSSSVNYSESNSTDSTKSQH HSSTSNQETSDSEMEMEAEHYPNGVLGSMSTRIVNGAYKHEDLQTDESSMDDRHPR

靶点信息

RAN is a small GTPase involved in the assembly of microtubules to form mitotic spindles. The protein encoded by this gene is a cytoplasmic guanine nucleotide exchange factor (GEF) that binds beta-tubulin and has GEF activity toward RAN. The encoded protein plays a role in the formation of noncentrosomal micro ubules. In addition, this protein may be involved in the regulation of D(1) receptor signaling by protein ki ase C delta and protein kinase C gamma. Several transcript variants encoding different isoforms have be en found for this gene. [provided by RefSeq, Feb 2016]
--

基因ID	57610
基因名	RANBP10
Swiss	Q6VN20
别名	RANBP10

产品验证



Western blot analysis of RANBP10 expressed in U-87MG,HeLa,Mouse spleen using RANBP10 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn