

CCDC50 Rabbit pAb

货号: B21141

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	35kDa/56kDa
实测分子量	36kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	LO2,293T,U-87MG,Mouse lung,Mouse brain,Rat brain
细胞定位	cytoplasm,cytosol
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 10-147 of human CCDC 50 (NP_777568.1).
序列	KLPGVKEVCRDFAVLEDHTLAHSLQEQEIEHHLASNVQRNRLVQHDLQVAKQLQEEDLKAQAQLQKRYKDLEQQDCEIAQEIQEKLAIEAERRRIQEKKDEDIARLLQEKELQEEKKRKKHFPEFPATRAYADSYYYE

靶点信息

研究背景	This gene encodes a soluble, cytoplasmic, tyrosine-phosphorylated protein with multiple ubiquitin-interacting domains. Mutations in this gene cause nonsyndromic, postlingual, progressive sensorineural DFNA44 hearing loss. In mouse, the protein is expressed in the inner ear during development and postnatal maturation and associates with microtubule-based structures. This protein may also function as a negative regulator of NF-κB signaling and as an effector of epidermal growth factor (EGF)-mediated cell signaling. Alternative splicing results in multiple transcript variants encoding distinct isoforms. [provided by RefSeq, Oct 2008]
------	--

基因ID	152137
基因名	CCDC50
Swiss	Q8IVM0
别名	C3orf6;DFNA44;YMER;CCDC50

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn