

MSL1 Rabbit pAb

货号: B21147

产品信息

| | |
|-------|--|
| 反应 | Human,Mouse,Rat |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:2000 |
| 理论分子量 | |
| 实测分子量 | 80kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | K-562,Mouse testis,Rat testis,Rat brain |
| 细胞定位 | nuclear speck,nucleoplasm,nucleus |
| 纯化 | Affinity purification |

抗原信息

| | |
|------|---|
| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 210-300 of human MSL1 (NP_001012241). |
| 序列 | LAVPSWRDHSVEPLRDPNPSDLLENLDDSVFSKRHAKLEDEKRRKRWDIQRIREQRILQRLQLRMYKKKGIQESEPEVTS FFPEPDDVES |


靶点信息

| | |
|------|--|
| 研究背景 | Component of histone acetyltransferase complex responsible for the majority of histone H4 acetylation at 'Lys-16' (H4K16ac which is implicated in the formation of higher-order chromatin structure. Greatly enhances MSL2 E3 ubiquitin ligase activity, promoting monoubiquitination of histone H2B at 'Lys-34' (H2BK34Ub). This modification in turn stimulates histone H3 methylation at 'Lys-4' (H3K4me and 'Lys-79' (H3K79me) and leads to gene activation, including that of HOXA9 and MEIS1. In the MSL complex, acts as a scaffold to tether MSL3 and KAT8 together for enzymatic activity regulation. |
|------|--|

| | |
|-------|------------------|
| 基因ID | 339287 |
| 基因名 | MSL1 |
| Swiss | Q68DK7 |
| 别名 | MSL-1;hMSL1;MSL1 |

产品验证

Western blot analysis of MSL1 expressed in Mouse testis,Rat testis,K-562 using MSL1 Rabbit pAb at 1:100. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



实验步骤

访问官网浏览详情: www.ablybio.cn