

# NPM1 Rabbit pAb

货号: **B21164** 

## 产品信息

| 成应 Human,Mouse,Rat  宿主 Rabbit  克隆性 Polyclonal  機測反应  応用 WB IHC IF/ICC  推荐液度 WB: 1;500 - 1:1000 IHC: 1:100 - 1:500 IF/ICC: 1:50 - 1:200  理论分子量 28kDa/29kDa/32kDa  実測分子量 38KDa  形式 Liquid  保存条件 Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.  の職物 Unconjugated  即性対照 293T,Jurkat,HeLa,NIH/3T3,Mouse thymus,Mouse spleen   |       |   |
|--|-------|---|
| 克隆性       Polyclonal         後期反应       Image: Main of the polyclonal of the pol | 反应    | Human,Mouse,Rat   |
| 度剤   | 宿主    | Rabbit  |
| 应用 WB IHC IF/ICC  推荐來度 WB: 1:500 - 1:1000 IHC: 1:100 - 1:500 IF/ICC: 1:50 - 1:200  理论分子量 28kDa/29kDa/32kDa  実測分子量 38KDa  形式 Liquid  保存条件 Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.  (興取物 Unconjugated  阳性対照 293T,Jurkat,HeLa,NIH/3T3,Mouse thymus,Mouse spleen   | 克隆性   | Polyclonal  |
| # # # # # # # # # # # # # # # # # # #  | 预测反应  |   |
| IHC: 1:100 - 1:500 IF/ICC: 1:50 - 1:200  理论分子量 28kDa/29kDa/32kDa  实测分子量 38KDa  形式 Liquid  R在条件 Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.  (周联物 Unconjugated  P性对照 293T,Jurkat,HeLa,NIH/3T3,Mouse thymus,Mouse spleen   | 应用    | WB IHC IF/ICC   |
| 实测分子量 38KDa  形式 Liquid  保存条件 Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.  偶联物 Unconjugated  阳性对照 293T,Jurkat,HeLa,NIH/3T3,Mouse thymus,Mouse spleen   | 推荐浓度  | IHC: 1:100 - 1:500  |
| 形式 Liquid  保存条件 Store at -20℃. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.  個联物 Unconjugated  阳性对照 293T,Jurkat,HeLa,NIH/3T3,Mouse thymus,Mouse spleen  | 理论分子量 | 28kDa/29kDa/32kDa   |
| 保存条件 Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.  (周联物 Unconjugated  四性对照 293T,Jurkat,HeLa,NIH/3T3,Mouse thymus,Mouse spleen   | 实测分子量 | 38KDa   |
| Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.  | 形式    | Liquid  |
| 阳性对照 293T,Jurkat,HeLa,NIH/3T3,Mouse thymus,Mouse spleen  | 保存条件  |   |
|  | 偶联物   | Unconjugated  |
| 细胞定位 Cytoplasm,Nucleus,centrosome,cytoskeleton,microtubule organizing center,nucleolus,nucleoplasm   | 阳性对照  | 293T,Jurkat,HeLa,NIH/3T3,Mouse thymus,Mouse spleen  |
|  | 细胞定位  | Cytoplasm, Nucleus, centrosome, cytoskeleton, microtubule organizing center, nucleolus, nucleoplasm |
| 纯化 Affinity purification   | 纯化    | Affinity purification   |

## 抗原信息


| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 9-122 of human NPM1 ( NP_002511.1).      |  |
|------|--|--|
| 序列   | MSPLRPQNYLFGCELKADKDYHFKVDNDENEHQLSLRTVSLGAGAKDELHIVEAEAMNYEGSPIKVTLATLKMSVQPTVSL<br>GGFEITPPVVLRLKCGSGPVHISGQHLVAVEED |  |

## 靶点信息

| 研究背景 | This gene encodes a phosphoprotein which moves between the nucleus and the cytoplasm. The gene pro duct is thought to be involved in several processes including regulation of the ARF/p53 pathway. A numb er of genes are fusion partners have been characterized, in particular the anaplastic lymphoma kinase ge ne on chromosome 2. Mutations in this gene are associated with acute myeloid leukemia. More than a do zen pseudogenes of this gene have been identified. Alternative splicing results in multiple transcript varia nts. |
|------|---|
|------|---|

| 基因 <b>ID</b> | 4869         |
|--------------|--------------|
| 基因名          | NPM1         |
| Swiss        | P06748       |
| 别名           | B23;NPM;NPM1 |

## 产品验证



Western blot analysis of NPM1 expressed in Mouse thymus,293T,Jurkat using NPM1 Rabbit pAb at 1:100 0. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Bloc king buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: www.ablybio.cn