

BMPR2 Rabbit pAb

货号: B21271

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	59kDa/115kDa
实测分子量	100kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa
细胞定位	Cell membrane,Single-pass type I membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 27-150 of human BMPR 2 (NP_001195.2).
序列	SQNQERLCAFKDPYQQDLGIGESRISHENGTILCSKGSTCYGLWEKSKGDINLVKQGCWSHIGDPQECHYEECVVTTTPP SIQNGTYRFCCCSTDLCNVNFTENFPPPDTTPLSPPHSFNRDET

靶点信息

研究背景	This gene encodes a member of the bone morphogenetic protein (BMP) receptor family of transmembran e serine/threonine kinases. The ligands of this receptor are BMPs, which are members of the TGF-beta su perfamily. BMPs are involved in endochondral bone formation and embryogenesis. These proteins transd uce their signals through the formation of heteromeric complexes of two different types of serine (threoni ne) kinase receptors: type I receptors of about 50-55 kD and type II receptors of about 70-80 kD. Type II r eceptors bind ligands in the absence of type I receptors, but they require their respective type I receptors for signaling, whereas type I receptors require their respective type II receptors for ligand binding. Mutati ons in this gene have been associated with primary pulmonary hypertension, both familial and fenflurami ne-associated, and with pulmonary venoocclusive disease.
基因 ID	659
基因名	BMPR2
Swiss	Q13873
别名	BMPR2;BMPR-II;BMPR3;BMR2;BRK-3;POVD1;PPH1;T-ALK

产品验证



Western blot analysis of BMPR2 expressed in HeLa using BMPR2 Rabbit pAb at 1:1000. Secondary antibod y: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn