

# TBC1D4 Rabbit pAb

货号: B21310

## 产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	146kDa
实测分子量	160kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Rat skeletal muscle
细胞定位	cytosol
纯化	Affinity purification


## 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1059-1298 of human TB C1D4 (NP_055647.2).
序列	SRLLHDYHRDLYNHLEENEISPSLYAAPWFLTLFASQFSLGFVARVFDIIFLQGTEVIFKVALSLLSSQETLIMECESFENIVEF LKNTLPDMNTSEMEKIITQVFEMDISKQLHAYEVEYHVLQDELQESSYSCEDSETLEKLERANSQLKRQNMDLLEKLQVAH TKIQALESNLENLLTRETKMKSLIRTLEQEKMAYQKTVEQLRKLLPADALVNCDLLLRDLNCNPNNKAKIGNKP

靶点信息

研究背景	This gene is a member of the Tre-2/BUB2/CDC16 domain family. The protein encoded by this gene is a Ra b-GTPase-activating protein, and contains two phopshotyrosine-binding domains (PTB1 and PTB2), a calm odulin-binding domain (CBD), a Rab-GTPase domain, and multiple AKT phosphomotifs. This protein is tho ught to play an important role in glucose homeostasis by regulating the insulin-dependent trafficking of the glucose transporter 4 (GLUT4), important for removing glucose from the bloodstream into skeletal muscle and fat tissues. Reduced expression of this gene results in an increase in GLUT4 levels at the plasma membrane, suggesting that this protein is important in intracellular retention of GLUT4 under basal conditions. When exposed to insulin, this protein is phosphorylated, dissociates from GLUT4 vesicles, resulting in increased GLUT4 at the cell surface, and enhanced glucose transport. Phosphorylation of this protein by AKT is required for proper translocation of GLUT4 to the cell surface. Individuals homozygous for a mutation in this gene are at higher risk for type 2 diabetes and have higher levels of circulating glucose and insulin levels after glucose ingestion. Alternative splicing results in multiple transcript variants encoding different isoforms.
基因ID	9882
基因名	TBC1D4
Swiss	O60343
别名	AS160;NIDDM5;TBC1D4

### 产品验证



Western blot analysis of TBC1D4 expressed in Rat skeletal muscle using TBC1D4 Rabbit pAb at 1:1000. Se condary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking b uffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

## 实验步骤

访问官网浏览详情: www.ablybio.cn