

SRGAP2 Rabbit pAb

货号: B21319

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	120kDa
实测分子量	140KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse brain
细胞定位	Cell junction,Cell membrane,Cell projection,Cytoplasm,Cytoplasmic vesicle,Nucleus,dendritic spine,lamelli podium,phagosome,postsynaptic cell membrane,postsynaptic density,synapse
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 790-900 of human SRG AP2 (NP_056141.2).
序列	GVVERSSPKSEIEVISEPPEEKVTARAGASCPSGGHVADIYLANINKQRKRPEGSIRKTFRSDSHGLSSLTDSSSPGVGA SCRPSQPIMSQLPKEGPDKCSISGHGS

靶点信息

研究背景	This locus encodes a member of the SLIT-ROBO Rho GTPase activating protein family. The encoded protein stimulates GTPase activity of Rac1, and plays a role in cortical neuron development. This locus has several paralogs on human chromosome 1 resulting from segmental duplication. While this locus itself is conserved among various species, the paralogs are found only in the genus Homo, and not in the genomes of non-human great apes. Alternatively spliced transcript variants have been described for this locus.
------	--

基因ID	23380
基因名	SRGAP2
Swiss	O75044
别名	SRGAP2;ARHGAP34;FNBP2;SRGAP2A;SRGAP3

产品验证



Western blot analysis of SRGAP2 expressed in Mouse brain using SRGAP2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn