

DNAJC9 Rabbit pAb

货号: B21475

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	29kDa
实测分子量	33KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	Cell membrane,Cytoplasm,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-260 of human DNAJC9 (NP_056005.1).
序列	MGLLDLCEEVFGTADLYRVLGVRREASDGEVRRGYHKVSLQVHPDRVGEGDKEDATRRFQILGKVYSVLSDREQRAVYD EQGTVDEDSPVLTQDRDWEAYWRLLFKKISLEDIQAFEKTYKGSEEELADIKQAYLDFKGDMDQIMESVLCVQYTEEPRIR NIIQQAIDAGEVPSYNAFVKESKQKMNARKRRAQEEAKEAEMSRKELGLDEGVDSLKAAIQSRQKDRQKEMDNFLAQME AKYCKSSKGGGKKSALKKEKK

靶点信息

研究背景	Acts as a dual histone chaperone and heat shock co-chaperone. As a histone chaperone, forms a co-chap erone complex with MCM2 and histone H3-H4 heterodimers; and may thereby assist MCM2 in histone H3-H4 heterodimer recognition and facilitate the assembly of histones into nucleosomes. May also act as a hi stone co-chaperone together with TONSL. May recruit histone chaperones ASF1A, NASP and SPT2 to histo ne H3-H4 heterodimers. Also plays a role as co-chaperone of the HSP70 family of molecular chaperone proteins, such as HSPA1A, HSPA1B and HSPA8. As a co-chaperone, may play a role in the recruitment of HS P70-type molecular chaperone machinery to histone H3-H4 substrates, thereby maintaining the histone st ructural integrity. Exhibits activity to assemble histones onto DNA in vitro.
基因 ID	23234
基因名	DNAJC9
Swiss	Q8WXX5
别名	DNAJC9;HDJC9;JDD1;SB73

产品验证



Western blot analysis of DNAJC9 expressed in 293T using DNAJC9 Rabbit pAb at 1:1000. Secondary antibo dy: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-f at dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn