

PSME4 Rabbit pAb

货号: B21484

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	24kDa/113kDa/139kDa/211kDa
实测分子量	211KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa
细胞定位	Cytoplasm,Nucleus,Nucleus speckle,cytosol
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1634-1843 of human PS ME4 (NP_055429.2).
序列	LYPHQVPLVLQVLKQTARSSSWHARYTVLTYLQTMVFYNLFIFLNNEDAVKDIRWLVISLLEDEQLEVREMAATTLSGLLQC NFLTMDSPMQIHFEQLCKTKLPKKRKRDPGSVGD TIPS AELVKRHAGVLGLGACVLSSPYDVPTWMPQLLMNLSAHLNDP QPIEMTVKKTLNFRRTHHDNWQE HKQQFTDDQLLVLTDLLVSPCYA

靶点信息

研究背景	Associated component of the proteasome that specifically recognizes acetylated histones and promotes ATP- and ubiquitin-independent degradation of core histones during spermatogenesis and DNA damage response. Recognizes and binds acetylated histones via its bromodomain-like (BRDL region and activates the proteasome by opening the gated channel for substrate entry. Binds to the core proteasome via its C-terminus, which occupies the same binding sites as the proteasomal ATPases, opening the closed structure of the proteasome via an active gating mechanism. Component of the spermatoproteasome, a form of the proteasome specifically found in testis: binds to acetylated histones and promotes degradation of histones, thereby participating actively to the exchange of histones during spermatogenesis. Also involved in DNA damage response in somatic cells, by promoting degradation of histones following DNA double-strand breaks.
基因ID	23198
基因名	PSME4
Swiss	Q14997
别名	PSME4;PA200

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn