

DNAJC2 Rabbit pAb

货号: B21555

产品信息

| 反应 | Human, Mouse |
|-------|---|
| | |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB |
| 推荐浓度 | WB: 1:500 - 1:2000 |
| 理论分子量 | 65kDa/71kDa |
| 实测分子量 | 80KDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | 293T |
| 细胞定位 | Cytoplasm,Nucleus,cytosol |
| 纯化 | Affinity purification |


抗原信息

| 抗原信息 | Recombinant fusion protein containing a sequence corresponding to amino acids 1-140 of human DNAJC2 (NP_055192.1). | |
|------|---|--|
| 序列 | MLLLPSAADGRGTAITHALTSASTLCQVEPVGRWFEAFVKRRNRNASASFQELEDKKELSEESEDEELQLEEFPMLKTLDP KDWKNQDHYAVLGLGHVRYKATQRQIKAAHKAMVLKHHPDKRKAAGEPIKEGDNDYFTC | |

靶点信息

| 研究背景 | This gene is a member of the M-phase phosphoprotein (MPP) family. The gene encodes a phosphoprotein with a J domain and a Myb DNA-binding domain which localizes to both the nucleus and the cytosol. The protein is capable of forming a heterodimeric complex that associates with ribosomes, acting as a molec ular chaperone for nascent polypeptide chains as they exit the ribosome. This protein was identified as a I eukemia-associated antigen and expression of the gene is upregulated in leukemic blasts. Also, chromos omal aberrations involving this gene are associated with primary head and neck squamous cell tumors. T his gene has a pseudogene on chromosome 6. Alternatively spliced variants which encode different protein isoforms have been described. |
|-------|--|
| 基因ID | 27000 |
| 基因名 | DNAJC2 |
| Swiss | Q99543 |
| 别名 | DNAJC2;MPHOSPH11;MPP11;ZRF1;ZUO1 |

产品验证



Western blot analysis of DNAJC2 expressed in 293T using DNAJC2 Rabbit pAb at 1:1000. Secondary antibo dy: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-f at dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn