

OX40L Rabbit pAb

货号: B21617

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	
实测分子量	30kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Raji,PC-3,HepG2,293T,Mouse thymus,Mouse spleen
细胞定位	cell surface,extracellular space,plasma membrane
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 51-130 of human OX40L (NP_003317.1).
序列	QVSHRYPRIQSIKVQFTEYKKEKGFIILTSQKEDEIMKVQNNSVIINCDFYLIISLKGYSQEVNISLHYQKDEEPLFQLK

靶点信息

研究背景	This gene encodes a cytokine of the tumor necrosis factor (TNF) ligand family. The encoded protein functions in T cell antigen-presenting cell (APC) interactions and mediates adhesion of activated T cells to endothelial cells. Polymorphisms in this gene have been associated with Sjogren's syndrome and systemic lupus erythematosus. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2014]
基因ID	7292

基因名	TNFSF4
Swiss	P23510
别名	CD134L;CD252;GP34;OX-40L;OX40L;TNLG2B;TXGP1;TNFSF4

产品验证



Western blot analysis of OX40L expressed in Mouse thymus,Raji,PC-3 using OX40L Rabbit pAb at 1:1000 . Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Block ing buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn