

SHANK2 Rabbit pAb

货号: B21633

产品信息

| 反应 | Human,Mouse,Rat |
|-------|---|
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | |
| 应用 | WB IF/ICC |
| 推荐浓度 | WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 |
| 理论分子量 | |
| 实测分子量 | 159kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | Mouse liver,Rat liver |
| 细胞定位 | apical plasma membrane, ciliary membrane, cytosol, dendritic spine, glutamatergic synapse, growth cone, hi ppocampal mossy fiber to CA3 synapse, neuron projection, plasma membrane, postsynaptic density, presyn aptic active zone |
| 纯化 | Affinity purification |

抗原信息

| 抗原信息 | A synthetic peptide corresponding to a sequence within amino acids 500-600 of human SHANK2 (NP_036 441.2). | |
|------|--|--|
| 序列 | PFALGANKDSLSAFEYPGPKRKLYSAVPGRLFVAVKPYQPQVDGEIPLHRGDRVKVLSIGEGGFWEGSARGHIGWFPAEC VEEVQCKPRDSQAETRADRSK | |

靶点信息

| 研究背景 | This gene encodes a protein that is a member of the Shank family of synaptic proteins that may function as molecular scaffolds in the postsynaptic density of excitatory synapses. Shank proteins contain multiple domains for protein-protein interaction, including ankyrin repeats, and an SH3 domain. This particular fa mily member contains a PDZ domain, a consensus sequence for cortactin SH3 domain-binding peptides a nd a sterile alpha motif. The alternative splicing demonstrated in Shank genes has been suggested as a mechanism for regulating the molecular structure of Shank and the spectrum of Shank-interacting protein s in the postsynaptic densities of the adult and developing brain. Alterations in the encoded protein may b e associated with susceptibility to autism spectrum disorder. Alternative splicing results in multiple transc ript variants. [provided by RefSeq, Feb 2014] |
|-------|---|
| 基因ID | 22941 |
| 基因名 | SHANK2 |
| Swiss | Q9UPX8 |
| 别名 | AUTS17;CORTBP1;CTTNBP1;ProSAP1;SHANK;SPANK-3;SHANK2 |

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn