

VDAC2 Rabbit pAb

货号: B22198

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:100 - 1:500 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	30kDa/31kDa/33kDa
实测分子量	35kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,HeLa,U-251MG,Mouse testis,Mouse brain,Mouse kidney,Rat brain,Rat kidney
细胞定位	Mitochondrion outer membrane
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human VDAC2 (NP_003366.2).
序列	MATHGQTCARPMCIPPSYADLGKAARDIFNKGFGFGLVKLDVKTKSCSGVEFSTSGSSNTDTGKVTGTLETKYKWCEYG LTFTEKWNTDNTLGTEIAIED

靶点信息

研究背景	This gene encodes a member of the voltage-dependent anion channel pore-forming family of proteins that are considered the main pathway for metabolite diffusion across the mitochondrial outer membrane. The encoded protein is also thought to be involved in the mitochondrial apoptotic pathway via regulation of BCL2-antagonist/killer 1 protein activity. Pseudogenes have been identified on chromosomes 1, 2, 12 and 21, and alternative splicing results in multiple transcript variants.
------	--

基因ID	7417
基因名	VDAC2
Swiss	P45880
别名	VDAC2; POR; voltage dependent anion channel 2

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn