

FAF2 Rabbit pAb

货号: B22694

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:1000 IF/ICC: 1:50 - 1:200
理论分子量	52kDa
实测分子量	53KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,293T,Mouse testis,Mouse brain,Rat liver
细胞定位	endoplasmic reticulum,extracellular region,lipid droplet
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 176-393 of human FAF2 (NP_055428.1).
序列	YLGHDDDHQDSDEFRCNTLCAPEVISLINRMLFWACSTNKPEGYRVSQALRENTYPFLAMIMLKDRRMTVVGRLLEGLIQPD DLINQLTFIMDANQTYLVSERLEREERNQTQVLRQQQDEAYLASLRADQEKERKKREERERKRRKEEEVQQQKLAEERRR QNLQEEKERKLECLPPEPSPDDPESVKIIFKLPNDSRVERRFHFSQSLTVIHDFLFS

靶点信息

研究背景	The protein encoded by this gene is highly expressed in peripheral blood of patients with atopi c (AD), compared to normal individuals. It may play a role in regulating the resistance to apoptosis that is observed in T cells and eosinophils of AD patients.
基因ID	23197

基因名	FAF2
Swiss	Q96CS3
别名	ETEA; UBXD8; UBXN3B

产品验证



Western blot analysis of FAF2 expressed in Mouse testis,Rat liver,HeLa using FAF2 Rabbit pAb at 1:1000 . Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Block ing buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn