

CDKN1A Rabbit pAb

货号: B22865

产品信息

反应	Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	17kDa
实测分子量	21KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse large intestine,Mouse spleen,Rat heart
细胞定位	cytoplasm,cytosol,nuclear body,nucleolus,nucleoplasm,nucleus,perinuclear region of cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-159 of mouse CDKN1 A (NP_031695.1).
序列	MSNPGDVRPVPHRSKVCRCCLFGPVDSEQLRRDCDALMAGCLQEARERWNFDFVTETPLEGNFVWERVRSGLPKVYLSP GSRSRDDLGGDKRPSTSSALLQGPAPEDHVALSLSCTLVSERPEDSPGGPGTSQGRKRRQTSLTDFYHSKRRLVFCKRK P

靶点信息

研究背景	This gene encodes a potent cyclin-dependent kinase inhibitor. The encoded protein binds to and inhibits the activity of cyclin-cyclin-dependent kinase2 or cyclin-dependent kinase4 complexes, and thus functions as a regulator of cell cycle progression at the G1 pahse. The expression of this gene is tightly controlled by the tumor suppressor protein p53, through which this protein mediates the p53-dependent cell cycle G1 phase arrest in response to a variety of stress stimuli. This protein can interact with proliferating cell nuclear antigen, a DNA polymerase accessory factor, and plays a regulatory role in S phase DNA replication and DNA damage repair. This protein was reported to be specifically cleaved by CASP3-like caspases, which thus leads to a dramatic activation of cyclin-dependent kinase2, and may be instrumental in the execution of apoptosis following caspase activation. Mice that lack this gene have the ability to regenerate damaged or missing tissue. Alternative splicing results in multiple transcript variants.
基因ID	12575
基因名	
Swiss	P39689
别名	CI; P2; SD; CAP; CDK; P21; Waf; mda; CDKI; CIP1; Cdkn; SDI1; Waf1; mda6; p21C; p21W; CAP20; Cdkn1; p21WAF; p21Cip1

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn