

MED26 Rabbit pAb

货号: B22881

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	65kDa
实测分子量	70KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Raji,Mouse brain,Mouse spinal cord
细胞定位	nucleoplasm,nucleus
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 500-574 of human MED 26 (NP_004822.2).
序列	SGAQTPGAHHFMSEYLKQEESTRQGARQLHVLVPQSPPTDLPGLTREVTQDDLDRIQASQWPGVNGCQDTQGNWY


靶点信息

研究背景
 切九月泉

The activation of gene transcription is a multistep process that is triggered by factors that recognize trans criptional enhancer sites in DNA. These factors work with co-activators to direct transcriptional initiation by the RNA polymerase II apparatus. The protein encoded by this gene is a subunit of the CRSP (cofactor required for SP1 activation) complex, which, along with TFIID, is required for efficient activation by SP1. This protein is also a component of other multisubunit complexes e.g. thyroid hormone receptor-(TR-) associated proteins which interact with TR and facilitate TR function on DNA templates in conjunction with initiation factors and cofactors.

基因ID	9441
基因名	MED26
Swiss	O95402
别名	CRSP7; CRSP70

产品验证



Western blot analysis of MED26 expressed in Raji,Mouse brain,Mouse spinal cord using MED26 Rabbit p Ab at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug p er lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn