

TIR1 Rabbit pAb

货号: B23452

产品信息

反应	Arabidopsis thaliana
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	66kDa
实测分子量	67KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Inflorescences
细胞定位	nucleus
纯化	Affinity purification

抗原信息


抗原信息	A synthetic peptide corresponding to a sequence within amino acids 495-594 of arabidopsis thaliana TIR1 (NP_567135.1).
序列	DKALLANASKLETMRSLWMSSCSVSFGACKLLGQKMPKLNVEIDERGAPDSRPESCPVERVFIYRTVAGPRFDMPGFV WNMDQDSTMRFSRQIITTNGL

靶点信息

研究背景	Encodes an auxin receptor that mediates auxin-regulated transcription. It contains leucine-rich repeats and an F-box and interacts with ASK1, ASK2 and AtCUL1 to form SCF-TIR1, an SCF ubiquitin ligase complex. Related to yeast Grr1p and human SKP2 proteins, involved in ubiquitin-mediated processes. Required for normal response to auxin and repressed in response to flagellin. As part of the SCF complex and in the presence of auxin, TIR1 interacts with Aux/IAA transcriptional repressor proteins and mediates their degradation. Mutations in TIR1 block auxin stimulation of flavonoid synthesis.
------	--

基因ID	825473
基因名	
Swiss	Q570C0
别名	AtTIR1; TRANSPORT INHIBITOR RESPONSE 1

产品验证



Western blot analysis of TIR1 expressed in Inflorescences using TIR1 Rabbit pAb at 1:1000. Secondary anti body: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% no n-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn