

# **DRP1** Rabbit pAb

货号: B23481

## 产品信息

反应	Human,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:100 - 1:500 IF/ICC: 1:50 - 1:200
理论分子量	60kDa/78kDa/79kDa/80kDa/81kDa/82kDa/83kDa
实测分子量	78-82KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Rat brain
细胞定位	Cytoplasm,Cytoplasmic vesicle,Endomembrane system,Golgi apparatus,Membrane,Mitochondrion outer membrane,Peripheral membrane protein,Peroxisome,clathrin-coated pit,cytosol,secretory vesicle,synaptic vesicle membrane
纯化	Affinity purification

## 抗原信息


抗原信息	Recombinant protein of human DRP1.
序列	

## 靶点信息

研究背景	This gene encodes a member of the dynamin superfamily of GTPases. The encoded protein mediates mito chondrial and peroxisomal division, and is involved in developmentally regulated apoptosis and program med necrosis. Dysfunction of this gene is implicated in several neurological disorders, including Alzheimer 's disease. Mutations in this gene are associated with the autosomal dominant disorder, encephalopathy, I ethal, due to defective mitochondrial and peroxisomal fission (EMPF). Alternative splicing results in multip le transcript variants encoding different isoforms.

基因ID	10059
基因名	DNM1L
Swiss	O00429
别名	DNM1L; DLP1; DRP1; DVLP; DYMPLE; EMPF; EMPF1; HDYNIV; dynamin 1 like

## 产品验证



Western blot analysis of DRP1 expressed in Rat brain using DRP1 Rabbit pAb at 1:1000. Secondary antibo dy: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-f at dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

### 实验步骤

访问官网浏览详情: www.ablybio.cn