

FH Rabbit pAb

货号**: B23521**

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	50kDa/54kDa
实测分子量	49KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	Cytoplasm,Mitochondrion
纯化	Affinity purification

抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 50-150 of human FH (NP_000134.2).	
序列	FRIEYDTFGELKVPNDKYYGAQTVRSTMNFKIGGVTERMPTPVIKAFGILKRAAAEVNQDYGLDPKIANAIMKAADEVAEGK LNDHFPLVVWQTGSGTQTN	

靶点信息

研究背景	The protein encoded by this gene is an enzymatic component of the tricarboxylic acid (TCA) cycle, or Kre bs cycle, and catalyzes the formation of L-malate from fumarate. It exists in both a cytosolic form and an N-terminal extended form, differing only in the translation start site used. The N-terminal extended form i s targeted to the mitochondrion, where the removal of the extension generates the same form as in the c ytoplasm. It is similar to some thermostable class II fumarases and functions as a homotetramer. Mutatio ns in this gene can cause fumarase deficiency and lead to progressive encephalopathy.
基因ID	2271

基因名	FH
Swiss	P07954
别名	FH; FMRD; HLRCC; LRCC; MCL; MCUL1; fumarate hydratase

产品验证



Western blot analysis of FH expressed in 293T using FH Rabbit pAb at 1:1000. Secondary antibody: HRP G oat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn