

Rig-I/DDX58 Rabbit pAb

货号: B23528

产品信息

反应	Human,Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	101kDa/106kDa
实测分子量	101kDa/106kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Jurkat,HepG2,MCF7
细胞定位	Cell junction,Cell projection,Cytoplasm,cytoskeleton,ruffle membrane,tight junction
纯化	Affinity purification

抗原信息


抗原信息	Recombinant protein of human Rig-I/DDX58.
序列	

靶点信息

研究背景	DEAD box proteins, characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD), are putative RNA helicases which are implicated in a number of cellular processes involving RNA binding and alteration of RNA secondary structure. This gene encodes a protein containing RNA helicase-DEAD box protein motifs and a caspase recruitment domain (CARD). It is involved in viral double-stranded (ds) RNA recognition and the regulation of immune response.
基因ID	23586

基因名	DDX58
Swiss	O95786
别名	DDX58; RIG-I; RIGI; RLR-1; SGMRT2; DExD/H-box helicase 58

产品验证



Western blot analysis of RIG-I/DDX58 expressed in Jurkat,HepG2,MCF7 using RIG-I/DDX58 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn