

MPP2 Rabbit pAb

货号: B23564

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	46kDa/60kDa/61kDa/63kDa/64kDa/66kDa
实测分子量	46kDa/60kDa/61kDa/63kDa/64kDa/66kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide, 50% glycerol, pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	cytoplasm, cytoskeleton, dendrite membrane, dendritic shaft, dendritic spine, plasma membrane, postsynaptic density
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-230 of human MPP2 (NP_005365.3).
序列	MPVAATNSETAMQQVLDNLGSLPSATGAAELDLIFLRGIMESPIVRSLAKAHERLEETKLEAVRDNNLELVQEILRDLAHVAEQSSTAAELAHILQEPHFQSLLETHDSVASKTYETPPPSPGLDPTFSNQPVPPDAVRMVGIRKTAGEHLGVTFREGGELVIRILHGGMVAQQQLLHVGDIIKEVNGQPVGSQDPRALQELLRNASGSVILKILPSYQEPHLPRQVFV

靶点信息

研究背景	Palmitoylated membrane protein 2 is a member of a family of membrane-associated proteins termed MAGUKs (membrane-associated guanylate kinase homologs). MAGUKs interact with the cytoskeleton and regulate cell proliferation, signaling pathways, and intracellular junctions. Palmitoylated membrane protein 2 contains a conserved sequence, called the SH3 (src homology 3) motif, found in several other proteins that associate with the cytoskeleton and are suspected to play important roles in signal transduction.
------	---

基因ID	4355
基因名	MPP2
Swiss	Q14168
别名	MPP2; DLG2; MAGUK p55 subfamily member 2

产品验证



Western blot analysis of MPP2 expressed in human esophagus cancer using MPP2 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn