

NRF2 Rabbit pAb

货号: B23578

产品信息

Human,Mouse,Rat
Rabbit
Polyclonal
WB IHC IF/ICC
WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
65kDa/66kDa/67kDa
97-100KDa
Liquid
Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
Unconjugated
C6
Cytoplasm,Nucleus,cytosol
Affinity purification

抗原信息


抗原信息	Recombinant protein of human NRF2.
序列	

靶点信息

研究背景	This gene encodes a transcription factor which is a member of a small family of basic leucine zipper (bZIP) proteins. The encoded transcription factor regulates genes which contain antioxidant response elements (ARE) in their promoters; many of these genes encode proteins involved in response to injury and inflam mation which includes the production of free radicals. Multiple transcript variants encoding different isofo rms have been characterized for this gene.
基因ID	4780

基因名	NFE2L2
Swiss	Q16236
别名	NFE2L2; HEBP1; NRF2; nuclear factor; erythroid 2 like 2

产品验证



Western blot analysis of NRF2 expressed in C6 using NRF2 Rabbit pAb at 1:1000. Secondary antibody: HR P Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn