

p53 Rabbit pAb

货号**: B23634**

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IHC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IP: 1:1000 - 1:5000
理论分子量	23-43kDa
实测分子量	23-43kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A431 cells, HEK-293 cells
细胞定位	Cytoplasm,Endoplasmic reticulum,Mitochondrion matrix,Nucleus,PML body
纯化	Affinity purification

抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 1-100 of human p53 (NP_000537.3).	
序列	MEEPQSDPSVEPPLSQETFSDLWKLLPENNVLSPLPSQAMDDLMLSPDDIEQWFTEDPGPDEAPRMPEAAPPVAPAPAAP TPAAPAPAPSWPLSSSVPSQ	

靶点信息

研究背景	This gene encodes a tumor suppressor protein containing transcriptional activation, DNA binding, and olig omerization domains. The encoded protein responds to diverse cellular stresses to regulate expression of target genes, thereby inducing cell cycle arrest, apoptosis, senescence, DNA repair, or changes in metab olism. Mutations in this gene are associated with a variety of human cancers, including hereditary cancer s such as Li-Fraumeni syndrome. Alternative splicing of this gene and the use of alternate promoters resu It in multiple transcript variants and isoforms. Additional isoforms have also been shown to result from th e use of alternate translation initiation codons from identical transcript variants (PMIDs: 12032546, 20937 277).
基因ID	7157
基因名	ТР53
Swiss	P04637
别名	TP53; BCC7; LFS1; P53; TRP53; tumor protein p53

产品验证



Western blot analysis of p53 expressed in A431 cells,HEK-293 cells using p53 Rabbit pAb at 1:1000. Secon dary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffe r: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn