

CDK2AP1 Rabbit pAb

货号: B23643

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:2000
理论分子量	9kDa/12kDa
实测分子量	9kDa/12kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HEK-293 cells, HeLa cells
细胞定位	cytoplasm,cytosol,nucleoplasm,nucleus,perinuclear region of cytoplasm
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-115 of human CDK2A P1 (NP_004633.1).
序列	MSYKPNLAAHMPAAALNAAGSVHSPSTSMATSSQYRQLLSDYGPPSLGYTQGTGNSQVPQSKYAELLAIIIEELGKEIRPTY AGSKSAMERLKRGIHARGLVRECLAETERNARS

靶点信息

研究背景	The protein encoded by this gene is a cyclin-dependent kinase 2 (CDK2) -associated protein which is thought to negatively regulate CDK2 activity by sequestering monomeric CDK2, and targeting CDK2 for proteolysis. This protein was found to also interact with DNA polymerase alpha/primase and mediate the phosphorylation of the large p180 subunit, which suggests a regulatory role in DNA replication during the S-phase of the cell cycle. This protein also forms a core subunit of the nucleosome remodeling and histone deacetylation (NURD) complex that epigenetically regulates embryonic stem cell differentiation. This gene thus plays a role in both cell-cycle and epigenetic regulation. Alternative splicing results in multiple transcript variants encoding distinct isoforms.
基因ID	8099
基因名	CDK2AP1
Swiss	O14519
别名	CDK2AP1; DOC1; DORC1; ST19; doc-1; p12DOC-1; cyclin-dependent kinase 2-associated protein 1

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn