

FADD Rabbit pAb

货号: B23658

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	23kDa
实测分子量	27kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-431,Mouse small intestine,Rat small intestine
细胞定位	cytoplasm, cytosol, neuron projection, nucleus, plasma membrane, ripoptosome
纯化	Affinity purification

抗原信息


抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 1-208 of human FADD (Q13158).
序列	MDPFLVLLHSVSSSLSSSELTELKFCLGRVGKRKLERVQSGLDLFSMILLEQNDLEPGHTELLRELLASLRRHDLLRRVDDF EAGAAAGAAPGEEDLCAAFNVICDNVGKDWRRLARQLKVSDTKIDSIEDRYPRNLTERVRESLRWKNTKENATVAHLVG ALRSCQMNVLADLVQEQQARDLQNRSGAMSPMSWNSDASTSEAS

靶点信息

研究背景	The protein encoded by this gene is an adaptor molecule that interacts with various cell surface receptors and mediates cell apoptotic signals. Through its C-terminal death domain, this protein can be recruited by TNFRSF6/Fas-receptor, tumor necrosis factor receptor, TNFRSF25, and TNFSF10/TRAIL-receptor, and thus it participates in the death signaling initiated by these receptors. Interaction of this protein with the receptors unmasks the N-terminal effector domain of this protein, which allows it to recruit caspase-8, and thereby activate the cysteine protease cascade. Knockout studies in mice also suggest the importance of this protein in early T cell development.
基因ID	8772
基因名	FADD
Swiss	Q13158
别名	FADD; GIG3; MORT1; Fas associated via death domain

产品验证

Western blot analysis of FADD expressed in A-431, Mouse small intestine, Rat small intestine using FADD Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



实验步骤

访问官网浏览详情: www.ablybio.cn