

MUC1 Rabbit pAb

货号: B23673

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	10-29kDa/103kDa/121-122kDa
实测分子量	10-29kDa/103kDa/121-122kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	
细胞定位	Apical cell membrane,Cell membrane,Cytoplasm,Nucleus,Secreted,Single-pass type I membrane protein
纯化	Affinity purification


抗原信息

抗原信息	A synthetic peptide corresponding to a sequence within amino acids 50-150 of human MUC1 (P15941).
序列	EKNAVSMTSSVLSSHSPGSGSSTTQGQDVTLPATEPASGSAATWGQDVTSVPVTRPALGSTTPPAHDVTSAPDNKPAP GSTAPPAHGVTSAPDTRPAPGS

靶点信息

研究背景	This gene encodes a membrane-bound protein that is a member of the mucin family. Mucins are O-glycosylated proteins that play an essential role in forming protective mucous barriers on epithelial surfaces. These proteins also play a role in intracellular signaling. This protein is expressed on the apical surface of epithelial cells that line the mucosal surfaces of many different tissues including lung, breast stomach and pancreas. This protein is proteolytically cleaved into alpha and beta subunits that form a heterodimeric complex. The N-terminal alpha subunit functions in cell-adhesion and the C-terminal beta subunit is involved in cell signaling. Overexpression, aberrant intracellular localization, and changes in glycosylation of this protein have been associated with carcinomas. This gene is known to contain a highly polymorphic variable number tandem repeats (VNTR) domain. Alternate splicing results in multiple transcript variants.
基因ID	4582
基因名	MUC1
Swiss	P15941
别名	MUC1; ADMCKD; ADMCKD1; CA 15-3; CD227; EMA; H23AG; KL-6; MAM6; MCD; MCKD; MCKD1; MUC-1; MUC-1/SEC; MUC-1/X; MUC1/ZD; PEM; PEMT; PUM; CA15-3; mucin-1

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn