

Phospho-FGFR1-Y653/Y654 Rabbit pAb

货号: B23847

产品信息

反应	Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	WB: 1:1000 - 1:5000
理论分子量	6kDa/16kDa/33kDa/55-95kDa
实测分子量	120KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	NIH/3T3
细胞定位	cytoplasmic vesicle,cytosol,extracellular region,nucleus,plasma membrane
纯化	Affinity purification


抗原信息

抗原信息	A synthetic phosphorylated peptide around Y653 & Y654 of human Phospho-FGFR1-Y653/Y654 (NP_0755 98.2).
序列	IDYYKK

靶点信息

研究背景	The protein encoded by this gene is a member of the fibroblast growth factor receptor (FGFR) family, whe re amino acid sequence is highly conserved between members and throughout evolution. FGFR family members differ from one another in their ligand affinities and tissue distribution. A full-length representative protein consists of an extracellular region, composed of three immunoglobulin-like domains, a single hy drophobic membrane-spanning segment and a cytoplasmic tyrosine kinase domain. The extracellular por tion of the protein interacts with fibroblast growth factors, setting in motion a cascade of downstream signals, ultimately influencing mitogenesis and differentiation. This particular family member binds both acid ic and basic fibroblast growth factors and is involved in limb induction. Mutations in this gene have been associated with Pfeiffer syndrome, Jackson-Weiss syndrome, Antley-Bixler syndrome, osteoglophonic dysplasia, and autosomal dominant Kallmann syndrome 2. Chromosomal aberrations involving this gene are a ssociated with stem cell myeloproliferative disorder and stem cell leukemia lymphoma syndrome. Alterna tively spliced variants which encode different protein isoforms have been described; however, not all variants have been fully characterized.
基因 ID	2260
基因名	FGFR1
Swiss	P11362
别名	FGFR1;BFGFR;CD331;CEK;ECCL;FGFBR;FGFR-1;FLG;FLT-2;FLT2;HBGFR;HH2;HRTFDS;KAL2;N-SAM;OGD;bFGF-R-1

产品验证



Western blot analysis of Phospho-FGFR1-Y653/Y654 expressed in NIH/3T3 using Phospho-FGFR1-Y653/Y654 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn