

XBP1S Rabbit pAb

货号: B24262

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:100 - 1:500 IF/ICC: 1:50 - 1:200
理论分子量	40KDa
实测分子量	60KDa/55KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T,Rat lung
细胞定位	
纯化	Affinity purification


抗原信息

抗原信息	Recombinant protein of human XBP1S.
序列	

靶点信息

研究背景	This gene encodes a transcription factor that regulates MHC class II genes by binding to a promoter elem ent referred to as an X box. This gene product is a bZIP protein, which was also identified as a cellular transcription factor that binds to an enhancer in the promoter of the T cell leukemia virus type 1 promoter. It may increase expression of viral proteins by acting as the DNA binding partner of a viral transactivator. It has been found that upon accumulation of unfolded proteins in the endoplasmic reticulum (ER), the mRN A of this gene is processed to an active form by an unconventional splicing mechanism that is mediated by the endonuclease inositol-requiring enzyme 1 (IRE1). The resulting loss of 26 nt from the spliced mRNA causes a frame-shift and an isoform XBP1(S), which is the functionally active transcription factor. The isof orm encoded by the unspliced mRNA, XBP1(U), is constitutively expressed, and thought to function as a n egative feedback regulator of XBP1(S), which shuts off transcription of target genes during the recovery p hase of ER stress. A pseudogene of XBP1 has been identified and localized to chromosome 5.
基因 ID	7494
基因名	XBP1
Swiss	P17861-2
别名	XBP1;TREB-5;TREB5;XBP-1;XBP2;XBP1s;CBX1

产品验证



Western blot analysis of XBP1S expressed in 293T,Rat lung using XBP1S Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn