

# Phospho-Acetyl CoA Carboxylase-S79 Rabbit pAb

货号: B24818

#### 产品信息

反应	Mouse
宿主	Rabbit
克隆性	Polyclonal
预测反应	
应用	WB
推荐浓度	<b>WB:</b> 1:500 - 1:2000
理论分子量	265kDa
实测分子量	280KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	NIH/3T3
细胞定位	
纯化	Affinity purification

#### 抗原信息

抗原信息	A synthetic phosphorylated peptide around S79 of human Acetyl CoA Carboxylase.
序列	

#### 靶点信息

#### 研究背景

Acetyl-CoA carboxylase (ACC) is a complex multifunctional enzyme system. ACC is a biotin-containing en zyme which catalyzes the carboxylation of acetyl-CoA to malonyl-CoA, the rate-limiting step in fatty acid synthesis. There are two ACC forms, alpha and beta, encoded by two different genes. ACC-alpha is highly enriched in lipogenic tissues. The enzyme is under long term control at the transcriptional and translation al levels and under short term regulation by the phosphorylation/dephosphorylation of targeted serine res idues and by allosteric transformation by citrate or palmitoyl-CoA. Multiple alternatively spliced transcript variants divergent in the 5' sequence and encoding distinct isoforms have been found for this gene.

基因ID	31,32
基因名	ACACA,ACACB
Swiss	Q13085,000763
别名	ACC; ACAC; ACC1; ACCA; Acac1; hACC1; ACACAD; ACCalpha; ACACalpha

## 产品验证



Western blot analysis of Phospho-Acetyl CoA Carboxylase-S79 expressed in NIH/3T3 using Phosph o-Acetyl CoA Carboxylase-S79 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit I gG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST . Detection: ECL Enhanced Kit. Exposure time: 120s.

### 实验步骤

访问官网浏览详情: www.ablybio.cn