

NF-κB p65/RelA Mouse mAb

货号: B25061

产品信息

反应	Human,Mouse,Rat
宿主	Mouse
克隆性	Monoclonal
预测反应	WB: Mononuclear macrophages , human cells , Mus musculus , Rattus norvegicus , Homo sapiens , Edw ardsiella tarda , Mauremys sinensis , Sus scrofa , Cyprinus carpio IF: human cells , Mus musculus , Rattus norvegicus , Homo sapiens IHC: Mus musculus , Sus scrofa , Rattus norvegicus Co-IP: Homo sapiens
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	58kDa/59kDa/60kDa
实测分子量	70kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Jurkat,HT-29,SGC-7901,NIH/3T3,Mouse liver
细胞定位	Cytoplasm,Nucleus
纯化	Affinity purification

抗原信息

抗原信息	Recombinant protein of human NF-κB p65
序列	

靶点信息

研究背景	NF-kappa-B is a ubiquitous transcription factor involved in several biological processes. It is held in the cytoplasm in an inactive state by specific inhibitors. Upon degradation of the inhibitor, NF-kappa-B moves to the nucleus and activates transcription of specific genes. NF-kappa-B is composed of NFKB1 or NFKB2 bound to either REL, RELA, or RELB. The most abundant form of NF-kappa-B is NFKB1 complexed with the product of this gene, RELA. Four transcript variants encoding different isoforms have been found for this gene.
基因ID	5970
基因名	RELA
Swiss	Q04206
别名	NFKB3;p65;NF-kB p65;RELA;CMCU

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn