

Phospho-ULK1-S555 Rabbit pAb

货号: B25085

产品信息

反应	Human
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Homo sapiens , Spodoptera frugiperda IHC: Homo sapiens
应用	WB
推荐浓度	WB: 1:500 - 1:1000
理论分子量	112kDa
实测分子量	113KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	Cytoplasm,Preautophagosomal structure,cytosol
纯化	Affinity purification


抗原信息

抗原信息	A synthetic phosphorylated peptide around S555 of human Phospho-ULK1-S555 (NP_003556.1).
序列	RLHSA

靶点信息

研究背景	Serine/threonine-protein kinase involved in autophagy in response to starvation. Acts upstream of phosphatidylinositol 3-kinase PIK3C3 to regulate the formation of autophagophores, the precursors of autophagosomes. Part of regulatory feedback loops in autophagy: acts both as a downstream effector and negative regulator of mammalian target of rapamycin complex 1 (mTORC1 via interaction with RPTOR. Activated via phosphorylation by AMPK and also acts as a regulator of AMPK by mediating phosphorylation of AMPK subunits PRKAA1, PRKAB2 and PRKAG1, leading to negatively regulate AMPK activity. May phosphorylate ATG13/KIAA0652 and RPTOR; however such data need additional evidences. Plays a role early in neuronal differentiation and is required for granule cell axon formation. May also phosphorylate SESN2 and SQSTM1 to regulate autophagy. Phosphorylates FLCN, promoting autophagy. Phosphorylates AMBRA1 in response to autophagy induction, releasing AMBRA1 from the cytoskeletal docking site to induce autophagosome nucleation.
基因ID	8408
基因名	ULK1
Swiss	O75385
别名	ULK1;ATG1;ATG1A;UNC51;Unc51.1;hATG1;ULK1

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn