

# TLR3 Rabbit pAb

货号: B25096

# 产品信息


反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Polyclonal
预测反应	WB: Mus musculus , Sus scrofa
应用	WB IHC IF/ICC
推荐浓度	WB: 1:100 - 1:500 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	72kDa/103kDa
实测分子量	104KDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.05% proclin300,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,Rat liver,Mouse lung
细胞定位	Early endosome,Endoplasmic reticulum membrane,Endosome membrane,Single-pass type I membrane pr otein
纯化	Affinity purification

# 抗原信息

抗原信息	Recombinant fusion protein containing a sequence corresponding to amino acids 500-700 of human TLR3 (NP_003256.1).
序列	SPFQPLRNLTILDLSNNNIANINDDMLEGLEKLEILDLQHNNLARLWKHANPGGPIYFLKGLSHLHILNLESNGFDEIPVEVFK DLFELKIIDLGLNNLNTLPASVFNNQVSLKSLNLQKNLITSVEKKVFGPAFRNLTELDMRFNPFDCTCESIAWFVNWINETHT NIPELSSHYLCNTPPHYHGFPVRLFDTSSCKDSA

研究背景	The protein encoded by this gene is a member of the Toll-like receptor (TLR) family which plays a fundam ental role in pathogen recognition and activation of innate immunity. TLRs are highly conserved from Dro sophila to humans and share structural and functional similarities. They recognize pathogen-associated molecular patterns (PAMPs) that are expressed on infectious agents, and mediate the production of cytok ines necessary for the development of effective immunity. The various TLRs exhibit different patterns of expression. This receptor is most abundantly expressed in placenta and pancreas, and is restricted to the dendritic subpopulation of the leukocytes. It recognizes dsRNA associated with viral infection, and induce s the activation of NF-kappaB and the production of type I interferons. It may thus play a role in host defe nse against viruses. Use of alternative polyadenylation sites to generate different length transcripts has b een noted for this gene.
基因ID	7098
基因名	TLR3
Swiss	O15455
别名	TLR3;CD283;IIAE2

# 产品验证



Western blot analysis of TLR3 expressed in HeLa,Rat liver,Mouse lung using TLR3 Rabbit pAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocki ng buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

# 实验步骤

访问官网浏览详情: www.ablybio.cn