

Rabbit anti Myc-Tag pAb

货号: B25152

产品信息

反应	
宿主	Rabbit
克隆性	Polyclonal
预测反应	Co-IP: Homo sapiens WB: Oryctolagus cuniculus , Solanum tuberosum , Arabidopsis thaliana , Homo sapiens , Mus musculus , Gallus gallus IHC: Mus musculus , Sus scrofa
应用	WB IF/ICC IP
推荐浓度	WB: 1:1000 - 1:10000 IF/ICC: 1:50 - 1:200 IP: 1:100 - 1:500
理论分子量	
实测分子量	
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	293T
细胞定位	
纯化	Affinity purification

抗原信息

抗原信息	A synthetic peptide corresponding to Myc tag.
序列	EQKLISEEDL

靶点信息

研究背景	Protein tags are peptide sequences genetically grafted onto a recombinant protein. Often these tags are removable by chemical agents or by enzymatic means, such as proteolysis or intein splicing. Tags are attached to proteins for various purposes. Epitope tags are short peptide sequences which are chosen because high-affinity antibodies can be reliably produced in many different species. These are usually derived from viral genes, which explain their high immunoreactivity. Epitope tags include V5-tag, Myc-tag, HA-tag and NE-tag. These tags are particularly useful for western blotting, immunofluorescence and immunoprecipitation experiments, although they also find use in antibody purification.
基因ID	
基因名	
Swiss	
别名	Myc;Myc tag;Myc-tag

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn