

Phospho-BRCA1-S988 Rabbit pAb

货号: **B25217**

产品信息

| | |
|-------|--|
| 反应 | Human |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | WB: Homo sapiens |
| 应用 | IF/ICC |
| 推荐浓度 | IF/ICC: 1:50 - 1:200 |
| 理论分子量 | 7kDa/78-85kDa/202-210kDa |
| 实测分子量 | Refer to figures |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.02% sodium azide,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | |
| 细胞定位 | Chromosome,Cytoplasm,Cytoplasm,Nucleus |
| 纯化 | Affinity purification |

抗原信息

| | |
|------|--|
| 抗原信息 | A synthetic phosphorylated peptide around S988 of human BRCA1 (NP_009225.1). |
| 序列 | IKSFV |

靶点信息

| | |
|-------|--|
| 研究背景 | This gene encodes a nuclear phosphoprotein that plays a role in maintaining genomic stability, and it also acts as a tumor suppressor. The encoded protein combines with other tumor suppressors, DNA damage sensors, and signal transducers to form a large multi-subunit protein complex known as the BRCA1-associated genome surveillance complex (BASC). This gene product associates with RNA polymerase II, and through the C-terminal domain, also interacts with histone deacetylase complexes. This protein thus plays a role in transcription, DNA repair of double-stranded breaks, and recombination. Mutations in this gene are responsible for approximately 40% of inherited breast cancers and more than 80% of inherited breast and ovarian cancers. Alternative splicing plays a role in modulating the subcellular localization and physiological function of this gene. Many alternatively spliced transcript variants, some of which are disease-associated mutations, have been described for this gene, but the full-length natures of only some of these variants has been described. A related pseudogene, which is also located on chromosome 17, has been identified. |
| 基因ID | 672 |
| 基因名 | BRCA1 |
| Swiss | P38398 |
| 别名 | BRCA1;BRCAI;BRCC1;BROVCA1;FANCS;IRIS;PNCA4;PPP1R53;PSCP;RNF53 |

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn