

Phospho-NF-κB2-S866 Rabbit pAb

货号: B25512

产品信息

| | |
|-------|---|
| 反应 | Human,Mouse |
| 宿主 | Rabbit |
| 克隆性 | Polyclonal |
| 预测反应 | IHC: Homo sapiens WB: Mus musculus , Rattus norvegicus , Capra hircus , Homo sapiens |
| 应用 | WB IHC IF/ICC |
| 推荐浓度 | WB: 1:500 - 1:2000 IHC: 1:50 - 1:100 IF/ICC: 1:50 - 1:100 |
| 理论分子量 | 47kDa/96kDa |
| 实测分子量 | 120kDa |
| 形式 | Liquid |
| 保存条件 | Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.01% thiomersal,50% glycerol,pH7.3. |
| 偶联物 | Unconjugated |
| 阳性对照 | ovary cancer |
| 细胞定位 | Cytoplasm,Nucleus |
| 纯化 | Affinity purification |

抗原信息

| | |
|------|--|
| 抗原信息 | A synthetic phosphorylated peptide around S866 of human NF-κB2 (NP_001070962.1). |
| 序列 | EDSAY |

靶点信息

| | |
|-------|--|
| 研究背景 | This gene encodes a subunit of the transcription factor complex nuclear factor-kappa-B (NFkB). The NFkB complex is expressed in numerous cell types and functions as a central activator of genes involved in inflammation and immune function. The protein encoded by this gene can function as both a transcriptional activator or repressor depending on its dimerization partner. The p100 full-length protein is co-translationally processed into a p52 active form. Chromosomal rearrangements and translocations of this locus have been observed in B cell lymphomas, some of which may result in the formation of fusion proteins. There is a pseudogene for this gene on chromosome 18. Alternative splicing results in multiple transcript variants. |
| 基因ID | 4791 |
| 基因名 | NFKB2 |
| Swiss | Q00653 |
| 别名 | CVID10;H2TF1;LYT-10;LYT10;NF-kB2;p100;p49/p100;p52 |

产品验证

实验步骤

访问官网浏览详情: www.ablybio.cn