


Tartaric acid disodium dihydrate

货号: **B26779**

产品信息

生物活性	Tartaric acid disodium dihydrate is a Acid phosphatase inhibitor , is a sodium salt used in buffers for molecular biology and cell culture applications. Increases the rate of colchicine binding to tubulin1.
CAS	6106-24-7
中文名称	
分子量	232.10
体外研究	
体内研究	
形式	Solid
运输条件	Room temperature in continental US; may vary elsewhere.
保存条件	4°C, sealed storage, away from moisture and light

溶解性	In Vitro: H₂O : 100 mg/mL (430.85 mM); Need ultrasonic) DMSO : (insoluble or slightly soluble) 配制储备液 <table><tr><th>浓度</th><th>溶剂</th><th>体积</th><th>质量</th><th>1 mg</th><th>5 mg</th><th>10 mg</th></tr><tr><td>1 mM</td><td>4.3085 mL</td><td>21.5424 mL</td><td>43.0849 mL</td><td></td><td></td><td></td></tr><tr><td>5 mM</td><td>0.8617 mL</td><td>4.3085 mL</td><td>8.6170 mL</td><td></td><td></td><td></td></tr><tr><td>10 mM</td><td>0.4308 mL</td><td>2.1542 mL</td><td>4.3085 mL</td><td></td><td></td><td></td></tr></table> * 请根据产品在不同溶剂中的溶解度选择合适的溶剂配制储备液；一旦配成溶液，请分装保存，避免反复冻融造成的产品失效。 储备液的保存方式和期限: -80℃, 6 months; -20℃, 1 month (sealed storage, away from moisture and light) 。-80℃ 储存时，请在 6 个月内使用，-20℃ 储存时，请在 1 个月内使用。 In Vivo: 请根据您的实验动物和给药方式选择适当的溶解方案。以下溶解方案都请先按照 In Vitro 方式配制澄清的储备液，再依次添加助溶剂： ——为保证实验结果的可靠性，澄清的储备液可以根据储存条件，适当保存；体内实验的工作液，建议您现用现配，当天使用； 以下溶剂前显示的百 分比是指该溶剂在您配制终溶液中的体积占比；如在配制过程中出现沉淀、析出现象，可以通过加热和/或超声的方式助溶 1. 请依序添加每种溶剂： PBS Solubility: 20 mg/mL (86.17 mM); Clear solution; Need ultrasonic *以上所有助溶剂都可在 MCE 网站选购。	浓度	溶剂	体积	质量	1 mg	5 mg	10 mg	1 mM	4.3085 mL	21.5424 mL	43.0849 mL				5 mM	0.8617 mL	4.3085 mL	8.6170 mL				10 mM	0.4308 mL	2.1542 mL	4.3085 mL			
浓度	溶剂	体积	质量	1 mg	5 mg	10 mg																							
1 mM	4.3085 mL	21.5424 mL	43.0849 mL																										
5 mM	0.8617 mL	4.3085 mL	8.6170 mL																										
10 mM	0.4308 mL	2.1542 mL	4.3085 mL																										
纯度	≥98.0%																												