

CCR9 Rabbit mAb

货号: B28411

产品信息

反应	Human, Mouse, Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB ICC FC
推荐浓度	WB: 1:500 - 1:2000 ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	40kDa/42kDa
实测分子量	45kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse brain
细胞定位	Cell membrane,Multi-pass membrane protein
纯化	Affinity purification
应用 推荐浓度 理论分子量 实测分子量 形式 保存条件 偶联物 阳性对照 细胞定位	WB: 1:500 - 1:2000 ICC: 1:50 - 1:200 FC: 1:20 - 1:50 40kDa/42kDa 45kDa Liquid Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3. Unconjugated Mouse brain Cell membrane,Multi-pass membrane protein


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human CCR9.
序列	QVTQTIAFFHSCLNPVLYVFVGERFRRDLVKTLKNLGCISQAQWVSFTRREGSLKLSSMLLETTSGALSL

靶点信息

研究背景	The protein encoded by this gene is a member of the beta chemokine receptor family. It is predicted to be a seven transmembrane protein similar to G protein-coupled receptors. Chemokines and their receptors are key regulators of the thymocytes migration and maturation in normal and inflammation conditions. The specific ligand of this receptor is CCL25. It has been found that this gene is differentially expressed by T lymphocytes of small intestine and colon, suggested a role in the thymocytes recruitment and develop ment that may permit functional specialization of immune responses in different segment of the gastroint estinal tract. This gene is mapped to the chemokine receptor gene cluster region. Two alternatively splice d transcript variants have been described. Multiple transcript variants encoding different isoforms have been found for this gene.
基因 ID	10803
基因名	CCR9
Swiss	P51686
别名	CCR9;CC-CKR-9;CDw199;GPR-9-6;GPR28

产品验证



Western blot analysis of CCR9 expressed in Mouse brain using CCR9 Rabbit mAb at 1:1000. Secondary ant ibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% no n-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn