

PKN2 Rabbit mAb

货号: B28453

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	75kDa/94kDa/106kDa/110kDa/112kDa
实测分子量	112kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Rat lung
细胞定位	apical junction complex,centrosome,cytoplasm,cytosol,intermediate filament cytoskeleton,nuclear body,n ucleoplasm,nucleus,perinuclear region of cytoplasm,plasma membrane
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human PKN2.
序列	GQDSETVFDIQNDRNSILPKSQSEYKPDTPQSGLEYSGIQELEDRRSQQRFQFNLQDFRCCAVLGRGHFGKVLLAEYKNT NEMFAIKALKKGDIVARDEVD

靶点信息

研究背景	PKC-related serine/threonine-protein kinase and Rho/Rac effector protein that participates in specific sign al transduction responses in the cell. Plays a role in the regulation of cell cycle progression, actin cytoskel eton assembly, cell migration, cell adhesion, tumor cell invasion and transcription activation signaling pr ocesses. Phosphorylates CTTN in hyaluronan-induced astrocytes and hence decreases CTTN ability to ass ociate with filamentous actin. Phosphorylates HDAC5, therefore lead to impair HDAC5 import. Direct Rho A target required for the regulation of the maturation of primordial junctions into apical junction formation in bronchial epithelial cells. Required for G2/M phases of the cell cycle progression and abscission during cytokinesis in a ECT2-dependent manner. Stimulates FYN kinase activity that is required for establishment of skin cell-cell adhesion during keratinocytes differentiation. Regulates epithelial bladder cells speed and direction of movement during cell migration and tumor cell invasion. Inhibits Akt pro-survival-induced kinase activity. Mediates Rho protein-induced transcriptional activation via the c-fos serum response factor (SRF. Involved in the negative regulation of ciliogenesis.
基因ID	5586
基因名	PKN2
Swiss	Q16513
别名	PAK2;PRKCL2;PRO2042;Pak-2;STK7;PKN2

产品验证



Western blot analysis of PKN2 expressed in Rat lung using PKN2 Rabbit mAb at 1:1000. Secondary antibod y: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn