

ATF2 Rabbit mAb

货号: B28460

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC ICC IP
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IC: 1:50 - 1:200 IP: 1:20 - 1:50
理论分子量	13- 15kDa/23-24kDa/35-54kDa
实测分子量	70kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HeLa,NIH/3T3
细胞定位	Cytoplasm,Mitochondrion outer membrane,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human ATF2.
序列	ТРТРТ

靶点信息

研究背景	This gene encodes a transcription factor that is a member of the leucine zipper family of DNA binding pro teins. The encoded protein has been identified as a moonlighting protein based on its ability to perform m echanistically distinct functions This protein binds to the cAMP-responsive element (CRE), an octameric p alindrome. It forms a homodimer or a heterodimer with c-Jun and stimulates CRE-dependent transcription . This protein is also a histone acetyltransferase (HAT) that specifically acetylates histones H2B and H4 in vitro; thus it may represent a class of sequence-specific factors that activate transcription by direct effect s on chromatin components. The encoded protein may also be involved in cell's DNA damage response in dependent of its role in transcriptional regulation. Several alternatively spliced transcript variants have b een found for this gene
基因ID	1386
基因名	ATF2
Swiss	P15336
别名	CRE-BP1;CREB-2;CREB2;HB16;TREB7;ATF2

产品验证



Western blot analysis of ATF2 expressed in HeLa,NIH/3T3 using ATF2 Rabbit mAb at 1:1000. Secondary an tibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% n on-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn