

INDO Rabbit mAb

货号: B28527

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200
理论分子量	45kDa
实测分子量	45kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	SKOV3,A-549
细胞定位	Cytoplasm,cytosol
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human INDO.
序列	MAHAMENSWTISKEYHIDEEVGFALPNPQENLPDFYNDWMFIAKHLPDLIESGQLRERVEKLNMLSIDHLTDHKSQRLARLVLGCITMAYVWGKGHDVRKVLPRNIAVPYCQLSKKLELPPILVYADCVLANWKKKDPNKPLTYENMDVLFsFRDGDCKSGFFLVSLLEIAAASAIKVIPTVFKAMQMQUERDTLLKALLEIASCLEKALQVFHQIHdHVNPKAFFSVLRIYLSGWKGNPQLSDGLVYEGFWEDPKFAGGSAGQSSVFQCFDVLGIIQQTAGGGHAAQFLQDMRRYMPPAHRNFLCSLESNPsvREFVLSKGDAGLREAYDACVKALVSLRSYHLQIVTKYILIPASQQPKENKTSedPSKLEAKGTGGTDLMNfLKTvrSTTEKSLLKEG

靶点信息

研究背景	This gene encodes indoleamine 2,3-dioxygenase (IDO) - a heme enzyme that catalyzes the first and rate-limiting step in tryptophan catabolism to N-formyl-kynurenine. This enzyme acts on multiple tryptophan substrates including D-tryptophan, L-tryptophan, 5-hydroxy-tryptophan, tryptamine, and serotonin. This enzyme is thought to play a role in a variety of pathophysiological processes such as antimicrobial and antitumor defense, neuropathology, immunoregulation, and antioxidant activity. Through its expression in dendritic cells, monocytes, and macrophages this enzyme modulates T-cell behavior by its pericellular catabolization of the essential amino acid tryptophan.
基因ID	3620
基因名	IDO1
Swiss	P14902
别名	IDO1;IDO;IDO-1;INDO;indoleamine 2

产品验证



实验步骤

访问官网浏览详情: www.ablybio.cn