

CCR3 Rabbit mAb

货号: B28537

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200
理论分子量	41kDa/43kDa
实测分子量	60kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	HepG2,A-549,Raji,Mouse liver,Rat spleen
细胞定位	Cell membrane,Multi-pass membrane protein
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human CCR3 .
序列	TVTFGVITSIVTWGLAVLAALPEFIFYETEELFEETLCSALYPEDTVYSWRHFHTLRMTIFCLVLPLLVMAICYTGIIKTL

靶点信息

研究背景	The protein encoded by this gene is a receptor for C-C type chemokines. It belongs to family 1 of the G pr otein-coupled receptors. This receptor binds and responds to a variety of chemokines, including eotaxin (CCL11), eotaxin-3 (CCL26), MCP-3 (CCL7), MCP-4 (CCL13), and RANTES (CCL5). It is highly expressed in e osinophils and basophils, and is also detected in TH1 and TH2 cells, as well as in airway epithelial cells. T his receptor may contribute to the accumulation and activation of eosinophils and other inflammatory cell s in the allergic airway. It is also known to be an entry co-receptor for HIV-1. This gene and seven other c hemokine receptor genes form a chemokine receptor gene cluster on the chromosomal region 3p21. Alter natively spliced transcript variants have been described.
基因ID	1232
基因名	CCR3
Swiss	P51677
别名	CCR3;CC-CKR-3;CD193;CKR3;CMKBR3

产品验证



Western blot analysis of CCR3 expressed in Mouse liver,Rat spleen,HepG2 using CCR3 Rabbit mAb at 1: 1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn