

ASK1 Rabbit mAb

货号: B28570

产品信息

反应	Human, Mouse
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC IF/ICC FC
推荐浓度	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200 IF/ICC: 1:50 - 1:200 FC: 1:20 - 1:50
理论分子量	69kDa/154kDa
实测分子量	155kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	A-431,HeLa
细胞定位	Cytoplasm,Endoplasmic reticulum
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human ASK1 .
序列	PPSPE

靶点信息

研究背景	Mitogen-activated protein kinase (MAPK) signaling cascades include MAPK or extracellular signal-regulate d kinase (ERK), MAPK kinase (MKK or MEK), and MAPK kinase kinase (MAPKKK or MEKK). MAPKK kinase/ME KK phosphorylates and activates its downstream protein kinase, MAPK kinase/MEK, which in turn activate s MAPK. The kinases of these signaling cascades are highly conserved, and homologs exist in yeast, Dros ophila, and mammalian cells. MAPKKK5 contains 1,374 amino acids with all 11 kinase subdomains. North ern blot analysis shows that MAPKKK5 transcript is abundantly expressed in human heart and pancreas. T he MAPKKK5 protein phosphorylates and activates MKK4 (aliases SERK1, MAPKK4) in vitro, and activates c-Jun N-terminal kinase (JNK)/stress-activated protein kinase (SAPK) during transient expression in COS and 293 cells; MAPKKK5 does not activate MAPK/ERK.
基因 ID	4217
基因名	MAP3K5
Swiss	Q99683
别名	MAP3K5;ASK1;MAPKKK5;MEKK5

产品验证



Western blot analysis of ASK1 expressed in A-431,HeLa using ASK1 Rabbit mAb at 1:1000. Secondary anti body: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. Blocking buffer: 5% no n-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn