

Myelin Basic Protein Rabbit mAb

货号: B28631

产品信息

反应	Human
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC
	WB: 1:500 - 1:2000 IHC: 1:50 - 1:200
理论分子量	18kDa
实测分子量	18kDa
形式	Liquid
	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Mouse brain,Rat brain
细胞定位 (Cytoplasmic side,Myelin membrane,Nucleus,Peripheral membrane protein
纯化 ,	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Myelin Basic Protein.
序列	GFLPRHRDTGILDSIGRFFGGDRGAPKRGSGKDSHHPARTAHYGSLPQKSHGRTQDENPVVHFFKNIVTPRTPPPSQGK GRGLSLSRFSWGAEGQRPGFGYGGRASDYKSAHKGFKGVDAQGTLSKIFKLGGRDSRSGSPMARR

靶点信息

研究背景	The protein encoded by the classic MBP gene is a major constituent of the myelin sheath of oligodendrocy tes and Schwann cells in the nervous system. However, MBP-related transcripts are also present in the bo ne marrow and the immune system. These mRNAs arise from the long MBP gene (otherwise called "Golli-MBP") that contains 3 additional exons located upstream of the classic MBP exons. Alternative splicing from the Golli and the MBP transcription start sites gives rise to 2 sets of MBP-related transcripts and gene p roducts. The Golli mRNAs contain 3 exons unique to Golli-MBP, spliced in-frame to 1 or more MBP exons. They encode hybrid proteins that have N-terminal Golli aa sequence linked to MBP aa sequence. The second family of transcripts contain only MBP exons and produce the well characterized myelin basic proteins. This complex gene structure is conserved among species suggesting that the MBP transcription unit is an integral part of the Golli transcription unit and that this arrangement is important for the function and/or regulation of these genes. [provided by RefSeq, Jul 2008]
基因ID	4155
基因名	MBP
Swiss	P02686
别名	MBP; myelin basic protein

产品验证



Western blot analysis of Myelin Basic Protein expressed in Mouse brain,Rat brain using Myelin Basic P rotein Rabbit mAb at 1:1000. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates /proteins: 30ug per lane. Blocking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. E xposure time: 120s.

实验步骤

访问官网浏览详情: www.ablybio.cn