

# Ikaros Rabbit mAb

货号: B28661

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IHC FC
推荐浓度	<b>WB:</b> 1:500 - 1:2000 <b>IHC:</b> 1:50 - 1:200 <b>FC:</b> 1:20 - 1:50
理论分子量	24-31kDa/41-57kDa
实测分子量	50-70kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	Raji
细胞定位	Cytoplasm,Nucleus,Nucleus
纯化	Affinity purification


抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Ikaros.
序列	EKENEMMKSHVMDQAINNAINYLGAESLRPLVQTTPPGGSEVVPVISPMYQLHKPLAEGTPRSNHSQAQDSAVENLLLLSKAK LVP SEREASPSNSCQDSTD

靶点信息

研究背景	This gene encodes a transcription factor that belongs to the family of zinc-finger DNA-binding proteins as associated with chromatin remodeling. The expression of this protein is restricted to the fetal and adult hemopoietic system, and it functions as a regulator of lymphocyte differentiation. Several alternatively spliced transcript variants encoding different isoforms have been described for this gene. Most isoforms share a common C-terminal domain, which contains two zinc finger motifs that are required for hetero- or homo-dimerization, and for interactions with other proteins. The isoforms, however, differ in the number of N-terminal zinc finger motifs that bind DNA and in nuclear localization signal presence, resulting in members with and without DNA-binding properties. Only a few isoforms contain the requisite three or more N-terminal zinc motifs that confer high affinity binding to a specific core DNA sequence element in the promoters of target genes. The non-DNA-binding isoforms are largely found in the cytoplasm, and are thought to function as dominant-negative factors. Overexpression of some dominant-negative isoforms have been associated with B-cell malignancies, such as acute lymphoblastic leukemia (ALL).
基因ID	10320
基因名	IKZF1
Swiss	Q13422
别名	IKZF1;CVID13;Hs.54452;IK1;IKAROS;LYF1;LyF-1;PPP1R92;PRO0758;ZNFN1A1

产品验证



实验步骤

访问官网浏览详情: [www.ablybio.cn](http://www.ablybio.cn)