

Nectin 2 Rabbit mAb

货号: B28668

产品信息

反应	Human,Mouse,Rat
宿主	Rabbit
克隆性	Monoclonal
预测反应	
应用	WB IF/ICC IP FC
推荐浓度	WB: 1:500 - 1:2000 IF/ICC: 1:50 - 1:200 IP: 1:20 - 1:50 FC: 1:20 - 1:50
理论分子量	51kDa/57kDa
实测分子量	62-75kDa
形式	Liquid
保存条件	Store at -20°C. Avoid freeze / thaw cycles. Buffer: PBS with 0.75% BSA,50% glycerol,pH7.3.
偶联物	Unconjugated
阳性对照	MCF-7,SW620,A-549,SKOV3
细胞定位	Cell membrane,Single-pass type I membrane protein
纯化	Affinity purification

抗原信息

抗原信息	Recombinant fusion protein corresponding to Human Nectin 2.
序列	FILLRVRRRKSPGGAGGGASGDGGFYDPKAQVLNGDPPVWTPVPGPMEPDGKDEEEEEEEKAEGKGLMLPPPALE DDMESQLDGS LISRRAVYV


靶点信息

研究背景	This gene encodes a single-pass type I membrane glycoprotein with two Ig-like C2-type domains and an Ig-like V-type domain. This protein is one of the plasma membrane components of adherens junctions. It also serves as an entry for certain mutant strains of herpes simplex virus and pseudorabies virus, and it is involved in cell to cell spreading of these viruses. Variations in this gene have been associated with differences in the severity of multiple sclerosis. Alternate transcriptional splice variants, encoding different isoforms, have been characterized.
------	-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

基因ID	5819
基因名	NECTIN2
Swiss	Q92692
别名	NECTIN2;CD112;HVEB;PRR2;PVRL2;PVRR2;nectin-2

产品验证

Western blot analysis of Nectin 2 expressed in MCF-7,SW620,A-549 using Nectin 2 Rabbit mAb at 1:100. Secondary antibody: HRP Goat Anti-Rabbit IgG (H+L) at 1:5000. Lysates/proteins: 30ug per lane. B locking buffer: 5% non-fat dry milk in TBST. Detection: ECL Enhanced Kit. Exposure time: 120s.



实验步骤

访问官网浏览详情: www.ablybio.cn